

Changes of Serum Ferritin in Acute Lung Injury Induced by Intestinal Ischemia/Reperfusion

Sung Dong Park¹ and Yoon-Yub Park²

¹Department of Chest Surgery, College of Medicine, Dongguk University, Gyeongju 780-714, ²Department of Physiology, School of Medicine, Catholic University of Daegu, Daegu 705-718, Korea

Serum ferritin levels are increased in subjects at-risk for or with acute lung injury (ALI), and there are observations to suggest that increases in serum ferritin levels may help predict the development of ALI in at-risk individuals. To deepen our understanding of increases of serum ferritin and their relationship to the development of ALI, we measured serum ferritin levels before and after intestinal ischemia/reperfusion (I/R) injury in rats, and found that serum ferritin levels increased significantly following I/R. Increases in serum and lavage ferritin levels paralleled increases in lung inflammation (lavage leukocyte numbers and tissue myeloperoxidase activities) and lung leak (lavage protein levels). In contrast, pre-treatment of rats with mepacrine (60 mg/kg, i.p.), a phospholipase A₂ inhibitor, attenuated not only I/R-induced serum and lavage ferritin increases, but also the development of ALI. These findings indicate that, besides of human subjects with ALI, serum ferritin levels increase early on also in an animal model of ALI. Therefore, serum and lavage ferritin can be a candidate for early biomarker of ALI.

Key Words: Intestinal ischemia/reperfusion, Acute lung injury, Ferritin, Rat

INTRODUCTION

Since Ashbaugh's early description of the acute respiratory distress syndrome (ARDS) in 1967 (Ashbaugh et al, 1967), morbidity and mortality have remained high for patients with ARDS (Ware & Matthay, 2000). ARDS usually occurs within 24 h of the insult and essentially occurs always within 72 h (Fowler et al, 1983; Hudson et al, 1995; Pepe et al, 1982), therefore, there is little time to enroll patients into studies.

Early and accurate identification of patients who are likely to develop ARDS would give several therapeutic advantages to the patients. Therefore, the development of specific biological markers for predicting ARDS is a desirable goal (Connelly & Repine, 1997). Numerous risk factors for the development of ARDS have been identified, and they are related to both direct and indirect causes of lung injury (Pittet et al, 1997). These markers may improve the prediction of ARDS in high risk clinical conditions and provide new insights into the pathogenesis of ARDS.

The iron storage protein, ferritin, plays a key role in iron metabolism. Its ability to sequester the element gives ferritin the dual functions of iron detoxification and iron reserve. Ferritin normally stores iron, but it may release free iron in the presence of superoxide or acidosis (Rief, 1992). For unknown reason, serum ferritin levels are increased in patients with acute lung injury (ALI) and mul-

ti-ple organ failure (MOF) (Connelly et al, 1997; Sharkey et al, 1999). Serum ferritin levels are also increased in patients at-risk for ALI development and appear to be increased more in at-risk patients who later develop ALI than in at-risk patients who do not develop ALI later. Additionally, serum ferritin levels correlated with the development of MOF (Sharkey et al, 1999), and severe hemorrhage induced ALI increases serum ferritin levels in rats (Park et al, 2003, 2004). However, the mechanisms responsible for the increase of serum ferritin and their relationship to ALI are still unclear. Therefore, the relationship between the pathogenesis of ALI and ferritin is worthy to be elucidated. Ischemia/reperfusion (I/R) occurs frequently in human pathological conditions (Grace, 1994), and intestinal I/R has been considered as a serious cause of ARDS (Otamiri et al, 1988).

In the present study, therefore, we evaluated serum ferritin level as a predictor of ALI in rats which were subjected to intestinal I/R injury. In addition, we examined the effect of pretreatment with mepacrine, a well-known phospholipase A₂ (PLA₂) inhibitor, on the changes of serum ferritin and ALI that occurs in I/R injured rats.

METHODS

Sprague-Dawley male rats, weighing 300~450 g, were used. Rats were anesthetized by ketamine (80 mg/kg, i.p.)

Corresponding to: Yoon-yub Park, Department of Physiology, School of Medicine, Catholic University of Daegu, 3056-6, Daemyeong-dong, Nam-gu, Daegu 705-718, Korea. (Tel) 82-53-650-4475, (Fax) 82-53-621-4106, (E-mail) yypark@cu.ac.kr

ABBREVIATIONS: ALI, acute lung injury; ARDS, acute respiratory distress syndrome; I/R, ischemia/reperfusion; MPO, myeloperoxidase.

and xylazine (16 mg/kg, i.p.). Left femoral artery was incised, and then polyethylene catheters (PE-50, Clay-Adams) filled with heparinized saline (100 U/ml) were inserted for blood sampling. The rats remained deeply anesthetized for the entire experiment, and the rats then were euthanized.

Rats were randomly divided into two groups and were given an intraperitoneal injection of PLA₂ inhibitor mepacrine (60 mg/kg, i.p.) or same volume of normal saline as a control. Thirty minutes after injection of saline or mepacrine, acute lung injury was induced by clamping superior mesenteric artery with a bulldog clamp for 30 min, and reperfusion was permitted for 3 hours. Sham-treated rats remained anesthetized after the same surgery without I/R.

For measurement of serum ferritin and protein concentrations, 100 µl of arterial blood was collected through the catheter before ischemia and 0, 30, 60, and 180 min after the reperfusion.

Bronchoalveolar lavage was performed by cannulating the trachea and instilling 8.0 ml of cold normal saline with a syringe. Saline was flushed three times through the lung. Approximately 6 ml of lavage fluid was recovered from each rat. Then, the fluid was centrifuged at 1,000 g for 10 min, and the supernatant was collected and stored at -70°C for additional assays. The cellular pellet was resuspended with 1.0 ml of distilled water and 1.0 ml of Hanks' balanced salt solution for a few seconds and centrifuged again. After supernatant was discarded, pellet was suspended in 0.4 ml of normal saline. Total leukocytes were counted with a hemocytometer.

Protein concentrations in serum and lavage fluid were measured using a bicinchoninic acid method (Sigma, St. Louis, MO). The ferritin content in serum and lavage fluid was quantified with commercially available ELISA kit specific for rat ferritin (Panapharm Laboratories, Kumamoto, Japan) by following the instructions.

To confirm pulmonary and intestinal infiltration of leukocytes, lung and intestinal tissues were collected and frozen at -70°C for analysis of myeloperoxidase (MPO) activity. MPO activity was measured by the method previously described (Lee et al, 2000). Briefly, the tissues were thawed

and then homogenized in 4 ml of potassium phosphate buffer (20 mM, pH 7.4), and the homogenate was centrifuged at 30,000 g for 30 min at 4°C. The pellet was resuspended in 4 ml of 50 mM potassium phosphate buffer (pH 6.0) containing 0.5% hexadecyltrimethylammonium bromide, sonicated for 90 s, and incubated for 2 h at 60°C. Then, MPO activity was determined using o-dianisidine as the substrate and H₂O₂ to initiate the reaction.

Statistical analyses: All data are presented as means±SEM. Unpaired t-test was used to compare groups, and paired t-test was used for paired observations within a group. A p value of <0.05 was considered statistically significant.

RESULTS

Effects of mepacrine on intestinal ischemia/reperfusion induced acute lung injury

Acute lung injury was induced in rats by means of intestinal I/R. The number of leukocyte (Fig. 1), protein content (Fig. 2) in lavage fluid, and lung MPO activity (Fig. 3) were significantly increased by intestinal I/R compared with those of Sham rats. As shown in Fig. 4, mepacrine treatment counteracted the increase of lavage leukocyte numbers, protein contents, and lung MPO activity. Intestinal MPO activity, however, did not show significant change after I/R.

Changes of serum and lung lavage ferritin levels after intestinal ischemia/reperfusion

Figure 5 shows the changes of serum ferritin levels in rats subjected to intestinal I/R. Thirty minutes of ischemia did not change serum ferritin levels. However, the values at 60 minutes after reperfusion showed sharp increases. Serum ferritin concentrations of I/R rats at 180 minutes following reperfusion were significantly higher than other groups. Mepacrine treatment significantly decreased serum ferritin increases in rats subjected to intestinal I/R. In




Fig. 1. Intestinal ischemia/reperfusion (I/R) significantly increased the number of leukocyte in bronchoalveolar lavage fluid. Mepacrine pretreatment (Mepa+I/R) effectively prevented the increase. The data shown are means±S.E.M. for the number of determinations shown in the parenthesis. *p<0.05, compared with Sham; #p<0.05, compared with I/R.




Fig. 2. Rats subjected to intestinal ischemia/reperfusion (I/R) had significantly increased protein contents in bronchoalveolar lavage fluid. This change was significantly decreased by the pretreatment with mepacrine (Mepa+I/R). The data shown are means±S.E.M. for the number of determinations shown in the parenthesis. *p<0.05, compared with Sham; ##p<0.01, compared with I/R.




Fig. 3. Intestinal ischemia/reperfusion (I/R) significantly elevated lung tissue myeloperoxidase (MPO) activity. This change was significantly attenuated by the pretreatment of rats with mepacrine (Mepa+I/R). The data shown are means±S.E.M. for the number of determinations shown in the parenthesis. ** $p < 0.01$, compared with Sham; # $p < 0.05$, compared with I/R.




Fig. 4. Intestinal tissue myeloperoxidase (MPO) activity did not change after intestinal ischemia/reperfusion (I/R) and mepacrine pretreated I/R (Mepa+I/R). The data shown are means±S.E.M. for the number of determinations shown in the parenthesis.

contrast, sham rats did not show any significant changes of serum ferritin throughout the experiment. Rats subjected to intestinal I/R had significantly increased lavage ferritin concentrations which were effectively decreased when rats were pretreated with mepacrine (Fig. 6).

DISCUSSION

Oxidative stress, defined as an imbalance caused by overproduction of free radicals or significant diminution in antioxidant defenses, is characterized by excessive levels of reactive oxygen species (ROS) that can cause cell injury and death in several organs (Halliwell, 1987). The reperfusion after severe but non-necrogenic ischemia generates ROS (Kirschner & Fantini, 1994). Intracellular free iron can catalyze the conversion of superoxide anion and hydrogen peroxide, produced by xanthine oxidase during reoxygenation, into more potent oxidants, such as hydroxyl




Fig. 5. Serum ferritin concentrations of Sham rats did not significantly change throughout the experiment. Rats subjected to intestinal ischemia/reperfusion (I/R) had greatly increased serum ferritin concentrations after 60 minutes of reperfusion. Serum ferritin concentrations of I/R rats at 180 minutes following reperfusion were significantly higher than those of other rats. By comparison, mepacrine pretreatment (Mepa+I/R) significantly attenuated the increase following I/R. Each point represents mean±S.E.M. for the number of determinations shown in parenthesis. ** $p < 0.01$, compared with Sham; # $p < 0.01$, compared with I/R.




Fig. 6. Intestinal ischemia/reperfusion (I/R) significantly increased bronchoalveolar lavage ferritin concentrations. Mepacrine pretreatment (Mepa+I/R) significantly counteracted the increase. The data shown are means±S.E.M. for the number of determinations shown in the parenthesis. * $p < 0.05$, compared with Sham; # $p < 0.05$, compared with I/R.

radicals (McCord, 1998), with deleterious effects on postischemic reperfused tissues. Indeed, several therapeutic strategies using iron chelation have proven to be useful to counteract the harmful effects of iron as a potentiator of I/R injury (Arora & Gores, 1996; Hybertson et al, 2002). Although the importance of iron in exacerbating the I/R damage is well recognized, the source of this catalytically active iron is not known. Ferritin, as an iron storage protein, can either represent a potentially harmful iron donor or an iron detoxicant (Harrison & Arosio, 1996).

In this study, mepacrine treatment was found to significantly reduce ALI in rats subjected to intestinal I/R and attenuate the increase of serum ferritin concentration. Park

et al. (2003, 2004) also reported recently that, in severe hemorrhage induced ALL, lung injury and increase of serum ferritin were attenuated in mepacrine-treated rats. These results confirm earlier study on the effects of mepacrine on intestinal I/R, intratracheal IL-1, and severe hemorrhage-induced lung injury (Lee et al, 1997, 1999; Park et al, 2004).

Serum ferritin started to increase at 60 min and reached about 10 times of basal values at 3 h after reperfusion. Mepacrine treatment significantly attenuated the increase of serum ferritin. Balla et al. (1992) also reported that the 1~2 h time lag was followed by rapid synthesis of ferritin in cultured endothelium which was exposed to heme.

Although serum ferritin levels correlate usually with total body iron stores, several physiological conditions such as inflammatory or infectious states (Harrison & Arosio, 1996) can increase it as one of the acute-phase proteins (Gabay & Kushner, 1999). Rogers et al. (1990) also reported that inflammatory induction of ferritin synthesis was different from iron dependent ferritin gene expression.

In general, hypoxia increases intracellular generation of ROS, and oxidative stress has been shown to mobilize iron from intracellular ferritin, thus increasing cell content of unbound iron and thereby promoting additional ferritin synthesis (Cairo et al, 1995; Chandel et al, 2000). Ferritin synthesis has been shown to increase cell resistance to oxidative stress by sequestering intracellular catalytic iron (Epsztejn et al, 1999; Garner et al, 1998), although various causes can elevate ferritin.

Serum ferritin levels may increase solely as a consequence of cellular necrosis or damage (Jacobs & Worwood, 1975). This possibility is supported by earlier clinical finding of an association between serum ferritin levels and injury severity score (Connelly et al, 1997; Sharkey et al, 1999). Furthermore, it is well established that proinflammatory systemic mediators, including TNF- α and IL-1, are elevated in intestinal I/R. Therefore, it is suggested that these inflammatory cytokines can increase the synthesis of ferritin (Rogers et al, 1990; Kwak et al, 1995) as one of the host defense mechanisms.

The importance of iron availability in amplifying I/R injury is well established. Elevated serum ferritin levels may also reflect an increase of ferritin synthesis as a protective response to increased oxidative stress, thereby increasing the ability to sequester iron and increasing antioxidant activity (Bolann & Ulvik, 1990; Balla et al, 1992, 1995; Vile & Tyrrell, 1993). Several studies also showed the protective effect of the iron chelator desferrioxamine (Arora & Gores, 1996; Hybertson et al, 2002). Although the origin of iron during in vivo postischemic reperfusion has not yet been identified, ferritin has been considered as a likely candidate source. Iron bound to ferritin is generally unavailable for the Harber-Weiss reaction, however, proteolysis, superoxide anion and acidosis, which are commonly encountered in disorders at risk for ARDS, may release iron from ferritin (Reif, 1992). Consequently, ferritin-derived iron may worsen oxidative damage in critically ill patients, contributing to the pathologic derangements seen in ARDS. Therefore, as described above, the exact role of increased serum ferritin is not clear at present, and additional studies may be required to explain these questions.

In conclusion, we found that serum ferritin levels were correlated with lung injury. Therefore, serum ferritin concentration in conjunction with other markers could be of

use in identifying individual patients at high risk of progressing to ARDS.

REFERENCES

- Arora AS, Gores GJ. The role of metals in ischemia/reperfusion injury of the liver. *Semin Liver Dis* 16: 31-38, 1996
- Ashbaugh DG, Bigelow DB, Petty TL, Levine BE. Acute respiratory distress in adults. *Lancet* 2: 319-323, 1967
- Balla G, Jacob HS, Balla J, Rosenberg M, Nath K, Apple F, Eaton JW, Vercellotti GM. Ferritin: a cytoprotective antioxidant strategem of endothelium. *J Biol Chem* 267: 18148-18153, 1992
- Balla J, Nath KA, Balla G, Juckett MB, Jacob HS, Vercellotti GM. Endothelial cell heme oxygenase and ferritin induction in rat lung by hemoglobin in vivo. *Am J Physiol* 268: L321-L327, 1995
- Bolann BJ, Ulvik RJ. On the limited ability of superoxide to release iron from ferritin. *Eur J Biochem* 193: 899-904, 1990
- Cairo G, Tacchini L, Pogliaghi G, Anzon E, Tomasi A, Bernelli-Zazzera A. Induction of ferritin synthesis by oxidative stress. Transcriptional and post-transcriptional regulation by expansion of the "free" iron pool. *J Biol Chem* 270: 700-703, 1995
- Chandel NS, Trzyna WC, McClintock DS, Schumacker PT. Role of oxidants in NF-kappa B activation and TNF-alpha gene transcription induced by hypoxia and endotoxin. *J Immunol* 165: 1013-1021, 2000
- Connelly KG, Moss M, Parsons PE, Moore EE, Moore FA, Giclas PC, Seligman PA, Repine JE. Serum ferritin as a predictor of the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 155: 21-25, 1997
- Connelly KG, Repine JE. Markers for predicting the development of acute respiratory distress syndrome. *Annu Rev Med* 48: 429-445, 1997
- Epsztejn S, Glickstein H, Picard V, Slotki IN, Breuer W, Beaumont C, Cabantchik ZI. H-ferritin subunit overexpression in erythroid cells reduces the oxidative stress response and induces multidrug resistance properties. *Blood* 94: 3593-3603, 1999
- Fowler AA, Hammon RF, Good JT, Benson KN, Baird M, Eberle DJ, Petty TL, Hyers TM. Adult respiratory distress syndrome: risk with common predisposition. *Ann Int Med* 98: 593-597, 1983
- Gabay C, Kushner I. Acute phase proteins and other systemic responses to inflammation. *N Engl J Med* 340: 448-454, 1999
- Garner B, Roberg K, Brunk UT. Endogenous ferritin protects cells with iron-laden lysosomes against oxidative stress. *Free Radic Res* 29: 103-114, 1998
- Grace PA. Ischaemia-reperfusion injury. *Br J Surg* 81: 637-647, 1994
- Halliwel B. Oxidants and human diseases: some new concepts. *FASEB J* 1: 358-364, 1987
- Harrison PM, Arosio P. The ferritins: molecular properties, iron storage function and cellular regulation. *Biochim Biophys Acta* 1275: 161-203, 1996
- Hudson LD, Milberg JA, Anardi D, Maunder RJ. Clinical risks for the development of the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 151: 293-301, 1995
- Hybertson BM, Connelly KG, Buser RT, Repine JE. Ferritin and desferrioxamine attenuate xanthine oxidase-dependent leak in isolated perfused rat lungs. *Inflammation* 26: 153-159, 2002
- Jacobs A, Worwood M. Ferritin in serum. Clinical and biochemical implications. *N Engl J Med* 292: 951-956, 1975
- Kirschner RE, Fantini GA. Role of iron and oxygen-derived free radicals in ischemia-reperfusion injury. *J Am Coll Surg* 179: 103-117, 1994
- Kwak EL, Larochelle DA, Beaumont C, Torti SV, Torti FM. Role of NF-kB in the regulation of ferritin H by tumor necrosis factor- α . *J Biol Chem* 270: 15285-15293, 1995
- Lee YM, Hybertson BM, Cho HG, Terada LS, Cho O, Repine AJ, Repine JE. Platelet activating factor contributes to acute lung leak in rats given interleukin-1 intratracheally. *Am J Physiol* 279: L75-80, 2000

- Lee YM, Hybertson BM, Terada LS, Repine AJ, Cho HG, Repine JE. Mepacrine decreases lung leak in rats given interleukin-1 intratracheally. *Am J Respir Crit Care Med* 155: 1624-1628, 1997
- Lee YM, Park Y, Kim T, Cho HG, Lee YJ, Repine JE. Effect of the inhibition of phospholipase A₂ in generation of free radicals in intestinal ischemia/reperfusion induced acute lung injury. *Korean J Physiol Pharmacol* 3: 263-273, 1999
- McCord JM. Iron, free radicals, and oxidative injury. *Semin Hematol* 35: 5-12, 1998
- Miller LL, Miller SC, Torti SV, Tsuji Y, Torti FM. Iron-independent induction of ferritin H chain by tumor necrosis factor. *Proc Natl Acad Sci USA* 88: 4946-4950, 1991
- Otamiri T, Lindahl M, Tagasson C. Phospholipase A₂ inhibition prevents mucosal damage associated with small intestine ischemia in rats. *Gut* 29: 489-494, 1988
- Park Y, Hybertson BM, Wright RM, Fini MA, Elkins ND, Repine JE. Serum ferritin elevation and acute lung injury in rats subjected to hemorrhage: reduction by mepacrine treatment. *Exp Lung Res* 30: 571-584, 2004
- Park YY, Hybertson BM, Wright RM, Repine JE. Serum ferritin increases in hemorrhaged rats that develop acute lung injury: effect of an iron-deficient diet. *Inflammation* 27: 257-263, 2003
- Pepe PE, Potkin RT, Reus DH, Hudson LD, Carrico CJ. Clinical predictors of the adult respiratory distress syndrome. *Am J Surg* 144: 124-130, 1982
- Pittet JF, Mackersie RC, Martin TR, Matthay MA. Biological markers of acute lung injury: prognostic and pathogenetic significance. *Am J Respir Crit Care Med* 155: 1187-1205, 1997
- Reif DW. Ferritin as a source of iron for oxidative damage. *Free Rad Biol Med* 12: 417-427, 1992
- Rogers JT, Bridges KR, Durmowicz GP, Glass J, Auron PE, Munro HN. Translational control during the acute phase response. Ferritin synthesis in response to interleukin-1. *J Biol Chem* 265: 14572-14578, 1990
- Sharkey RA, Donnelly SC, Connelly KG, Robertson CE, Haslett C, and Repine JE. Initial serum ferritin levels in patients with multiple trauma and the subsequent development of acute respiratory distress syndrome. *Am J Respir Crit Care Med* 159: 1506-1509, 1999
- Vile GF, Tyrrell RM. Oxidative stress resulting from ultraviolet A irradiation of human skin fibroblasts leads to a heme oxygenase-dependent increase in ferritin. *J Biol Chem* 268: 14678-14681, 1993
- Ware LB and Matthay MA. The acute respiratory syndrome. *N Engl J Med* 342: 1334-1349, 2000
-